

Heavy–light decay constants from the step scaling method*

Filippo Palombi^a

^a“E. Fermi” Research Center, c/o Compendio Viminale – pal. F, I-00184 Rome, Italy

We discuss results for the heavy–light decay constants in the continuum limit of quenched lattice QCD from finite size scaling techniques. We disentangle the simultaneous presence of the different energy scales characterizing heavy–light physics by first performing simulations at the unphysical volume $L_0 = 0.4$ fm, and then evolving the results towards the infinite volume. We find $f_{B_s} = 192(6)(4)$ MeV and $f_{D_s} = 240(5)(5)$ MeV. The approach has been developed by the APE group at the University of Rome “Tor Vergata”.

1. Introduction

The decay constants $f_{h\ell}$ of the heavy–light pseudoscalar mesons depend upon two different energy scales. A first energy scale is the one associated with the heavy quark mass m_h , and a second one is related to the light quark mass m_ℓ , or Λ_{QCD} . As a consequence, numerical simulations of heavy–light decay constants require tiny lattice spacings, in order to properly describe the dynamics of the heavy quark (highly localized), and large physical volumes, to accommodate the light quark (widely spread). In practice, a direct reliable simulation of $f_{h\ell}$ should demand a lattice size of around $L/a \sim O(100)$, which is hardly affordable nowadays in quenched simulations as well as in future unquenched ones. Different approaches are available on the market to face this problem [3]. I describe a new method which has been developed by the APE group at the University of Rome “Tor Vergata” [1,2], called the *step scaling method* at Lattice 2002 conference [4].

2. Step scaling method

The step scaling method is a two step procedure where the heavy–light decay constants $f_{h\ell}$ are first computed on a small unphysical volume with $L_0 = 0.4$ fm, and then evolved towards larger volumes in order to remove finite size effects. The evolution is simply provided by the

identity

$$f_{h\ell}(L_\infty) = f_{h\ell}(L_0) \frac{f_{h\ell}(L_1)}{f_{h\ell}(L_0)} \frac{f_{h\ell}(L_2)}{f_{h\ell}(L_1)} \dots \quad (1)$$

where $L_0 < L_1 < L_2 \dots$. The basic ingredient of the procedure is the computation of the step scaling function

$$\sigma(m_\ell, m_h, L_{k-1}) = \left. \frac{f_{h\ell}(m_\ell, m_h, L_k)}{f_{h\ell}(m_\ell, m_h, L_{k-1})} \right|_{L_k = s L_{k-1}} \quad (2)$$

i.e. the ratio of the decay constants at two different volumes and same quark masses. We fix $s = 2$ as the ratio of subsequent sizes and, after two evolution steps, we end up on $L_2 = 1.6$ fm, which we take as the “infinite” volume, i.e. free of finite size effects. At fixed lattice spacing, both $f_{h\ell}(L_0)$ and the σ ’s are simulated for a set of different quark masses, measured in a RGI quark mass scheme. Then, they are separately extrapolated to the continuum limit. Units are fixed through the r_0 scale [5,6]. The main advantage of the method is that $f_{h\ell}(L_0)$ can be simulated at physical heavy quark masses without big lattice artifacts while the σ ’s preserve in all steps a soft dependence upon the heavy quark mass, due to cancellations among volume depending terms. This makes the extrapolation to the physical values of m_c^{RGI} and m_b^{RGI} numerically safe. Moreover, such values can be obtained by a self-consistent application of the step scaling method to the HL–meson masses [7,8].

*talk given at Lattice 2003

3. Computational framework

The calculation is set up on the Schrödinger Functional scheme, with topological parameters

$$T = 2L, \quad C = C' = 0, \quad \theta = 0 \quad (3)$$

C and C' being the gauge fields on time boundaries, and θ being a phase which affects the periodicity of the fermion spatial boundary conditions. We simulate the basic SF correlation functions, i.e. the axial, pseudoscalar and boundary normalization bilinears

$$F_A^I(x_0) = -\frac{a^6}{2} \sum_{\mathbf{y}\mathbf{z}} \langle \bar{\zeta}_j(\mathbf{y}) \gamma_5 \zeta_i(\mathbf{z}) A_0^I(x) \rangle \quad (4)$$

$$F_P(x_0) = -\frac{a^6}{2} \sum_{\mathbf{y}\mathbf{z}} \langle \bar{\zeta}_j(\mathbf{y}) \gamma_5 \zeta_i(\mathbf{z}) P(x) \rangle \quad (5)$$

$$F_1 = -\frac{a^{12}}{3L^6} \sum_{\mathbf{y}\mathbf{z}\mathbf{y}'\mathbf{z}'} \langle \bar{\zeta}_j(\mathbf{y}) \gamma_5 \zeta_i(\mathbf{z}) \bar{\zeta}_i'(\mathbf{y}') \gamma_5 \zeta_j'(\mathbf{z}') \rangle \quad (6)$$

In terms of these, a finite size definition of the decay constants is given by taking the axial correlation function in the middle of the lattice

$$f_{h\ell} = \frac{2}{\sqrt{L^3 M_A(T/2)}} Z_A (1 + b_A a m) \frac{F_A^I(T/2)}{\sqrt{F_1}} \quad (7)$$

and the normalization is provided in terms of the effective meson mass, still measured in a finite size fashion

$$aM_A(T/2) = \frac{1}{2} \ln \left[\frac{F_A^I(T/2 - a)}{F_A^I(T/2 + a)} \right] \quad (8)$$

Quark flavors are defined according to their RGI masses (scale/finite-size independent), defined as follows. We start from the bare quark masses

$$am_i = \frac{1}{2} \left[\frac{1}{\kappa_i} - \frac{1}{\kappa_c} \right] \quad (9)$$

$$m_{ij}^{WI} = \frac{(1/2)(\partial_0 + \partial_0^*)F_A + ac_A \partial_0^* \partial_0 F_P}{2F_P} \Big|_{T/2} \quad (10)$$

and connect them to the RGI quark masses through proper renormalisation factors known in literature [9,10]

$$m_{ij}^{RGI} = Z_M \left[1 + (b_A - b_P) \frac{am_i + am_j}{2} \right] m_{ij}^{WI} \quad (11)$$

$$\hat{m}_i^{RGI} = Z_M Z [1 + b_m am_i] m_i, \quad Z = \frac{Z_m Z_P}{Z_A} \quad (12)$$

In this way, we get a plethora of definitions, all with the same continuum limit and differing in $O(a^2)$ terms:

1. $m_i^{RGI} = m_{ii}^{RGI}$
2. $m_{i(j)}^{RGI} = 2m_{ij}^{RGI} - m_{jj}^{RGI}, j \neq i$
3. $\hat{m}_i^{RGI}$

After choosing a target pair (m_i^{RGI}, m_j^{RGI}) , we tune the hopping parameters (κ_i, κ_j) such that the values of the masses coming from the various definitions equate the chosen pair. Of course, each choice of (κ_i, κ_j) produce a different value of $f_{h\ell}$, all the values differing in $O(a^2)$ terms, as it can be seen in Fig.1.




Figure 1. Continuum extrapolation on the small volume of $f_{B_s}(L_0)$. Units in GeV.

4. Results at $L_0 = 0.4$ fm

Simulations of the decay constants on the smallest volume $L_0 = 0.4$ fm have been performed at five different lattice spacings, using the geometries 24×12^3 , 32×16^3 , 40×20^3 , 48×24^3 and 64×32^3 . For each lattice spacing a set of eight RGI quark masses have been simulated, two around m_b^{RGI} , two around m_c^{RGI} , one additional at 4.0 GeV, and three light quark masses at 140, 100 and 60 MeV. We fix the light quark mass at the physical value of m_s^{RGI} by interpolation and we get

$$f_{B_s}(L_0) = 475(2) \text{ MeV} \quad (13)$$

$$f_{D_s}(L_0) = 644(3)\text{MeV} \quad (14)$$

The errors quoted at this stage are statistical only, and they have been evaluated by a jack-knife procedure. The continuum extrapolation has been obtained by performing a combined linear fit in $(a/r_0)^2$ over the last three sets of points of Fig.1. We estimate a systematic error of 1%, related to this extrapolation, by the discrepancy observed after restricting the fit to the two sets of points nearest to the continuum. The results obtained are clearly unphysical, as they have to be corrected through the evolution factors given by the σ 's.

5. Evolution steps

The first evolution step connects L_0 to $L_1 = 2L_0 = 0.8$ fm. Three different lattice spacings have been simulated in order to perform the continuum extrapolation, corresponding to the geometries with $L_0/a = 8, 12, 16$ and consequently $L_1/a = 16, 24, 32$. The values of the simulated quark masses have been halved in order to leave the lattice artifacts unchanged. In Fig. 2 the continuum step scaling function $\sigma(L_0)$ is plotted vs. the inverse of the heavy quark mass with the light quark mass fixed at m_s^{RGI} . At this stage we get




Figure 2. Continuum step scaling function $\sigma(L_0)$ as a function of $1/m_h^{RGI}$ at $m_\ell^{RGI} = m_s^{RGI}$. Units in GeV.

$$\sigma_{B_s}(L_0) = 0.417(3) \quad (15)$$

$$\sigma_{D_s}(L_0) = 0.414(3) \quad (16)$$

Analogously, the second evolution step connects L_1 to $L_2 = 2L_1 = 1.6$ fm. Also here, three lattice spacings have been simulated with the same lattice geometries as before. Again the values of the simulated quark masses have been halved. Fig. 3 shows the continuum step scaling function

$\sigma(L_1)$ vs. the inverse of the heavy quark mass at $m_\ell^{RGI} = m_s^{RGI}$. We find

$$\sigma_{B_s}(L_1) = 0.97(3) \quad (17)$$

$$\sigma_{D_s}(L_1) = 0.90(2) \quad (18)$$




Figure 3. Continuum step scaling function $\sigma(L_1)$ as a function of $1/m_h^{RGI}$ at $m_\ell^{RGI} = m_s^{RGI}$. Units in GeV.

6. Results

Combining the results of the small volume L_0 with the evolution factors, we obtain

$$f_{B_s} = 192(6)(4) \text{ MeV} \quad (19)$$

$$f_{D_s} = 240(5)(5) \text{ MeV} \quad (20)$$

The first error is statistical while the second one is systematic and due to the uncertainties on the continuum extrapolation, on the scale and on the renormalisation factors.

REFERENCES

1. M. Guagnelli et al., Phys. Lett. B546 (2002) 237, hep-lat/0206023.
2. G. M. de Divitiis, M. Guagnelli, F. Palombi, R. Petronzio and N. Tantalo, arXiv:hep-lat/0307005.
3. Andreas S. Kronfeld, these proceedings.
4. N. Yamada, arXiv:hep-lat/0210035.
5. S. Necco and R. Sommer, Nucl. Phys. B **622** (2002) 328
6. M. Guagnelli, R. Petronzio and N. Tantalo, Phys. Lett. B **548** (2002) 58
7. N. Tantalo, these proceedings.
8. G. M. de Divitiis, M. Guagnelli, R. Petronzio, N. Tantalo and F. Palombi, arXiv:hep-lat/0305018.
9. S. Capitani, M. Guagnelli, M. Luescher, S. Sint, R. Sommer, P. Weisz and H. Wittig, Nucl. Phys. Proc. Suppl. **63** (1998) 153
10. J. Rolf and S. Sint [ALPHA Collaboration], JHEP **0212** (2002) 007